arXiv:1610.00807v2 [math.NT] 13 Oct 2016

PERIODIC POINTS OF QUADRATIC POLYNOMIALS IN GALOIS EXTENSIONS

ROBIN ZHANG

Abstract. For quadratic polynomials with coefficients in a field k of characteristic zero, we seek to understand the periodic points under the actions of its iteration and the absolute Galois group $\mathrm{Gal}(\bar{k}/k)$. Supported by previous numerical work, we conjecture that these two actions are highly correlated. When $k = \mathbf{Q}$, we show that this conjecture implies the non-existence of quadratic periodic points of period 5 and prove that this conjecture is true for almost all rational c (in the sense of height density) using Hilbert's irreducibility theorem.

1. Introduction

The focus in the study of periodic points of quadratic polynomials (see Definition 2.1) has generally restricted both the coefficients and domain to the rational numbers. For instance, it has been demonstrated over the past two decades that there are no quadratic polynomials with rational fixed points of exact period 4, 5, or (conditionally) 6 [9, 6, 17]. In general, significant attention has centered on the conjecture of Poonen [14] that there are no such points with exact period greater than 3.

When the domain of quadratic polynomials with rational coefficients to quadratic number fields is lifted to quadratic number fields, there is no clear analogue to the non-existence conjecture of Poonen. Periodic points of quadratic polynomials with rational coefficients exist in quadratic number fields and those with exact period less than 5 have been well-studied (e.g. parametrization) due to several previous works [5, 9, 12, 13, 18]. However, the classification of periodic points of such quadratic polynomials with period ≥ 5 in quadratic number fields or higher degree extensions is still unknown. For instance, the existence of periodic points with exact period 5 in quadratic number fields is still an open conjecture:

Conjecture 1.1. There are no rational values c such that $\phi_c(z) = z^2 + c$ has a periodic point of exact period 5 that is quadratic over $\mathbf{Q}$.

Remark 1.2. Conjecture 1.1 is supported by a computational search on 150 million possible orbits on $C_0(5)$ (as described by Flynn, Poonen, and Schaefer [6]) that returned no positive results [19]. More extensive numerical evidence is provided by Hutz and Ingram [7] for the unconditional version of Theorem 1.6. Finiteness of the set of quadratic periodic points of ϕ_c with period 5 is well-known due to the high genus of the algebraic curves $C_1(5)$ and $C_0(5)$, as described by Flynn, Poonen, and Schaefer [6], and an appropriate application of Faltings's theorem.

Date: October 14, 2016.

Remark 1.3. Doyle [3] proved that Conjecture 1.1 holds for a quadratic field K when the Jacobian of the curve describing the 5-cycles has rank 1 over K. However, the Jacobian has greater rank for certain quadratic fields (e.g. $\mathbf{Q}(i)$), so this does not yield the full result. There also exists a local-global approach to Conjecture 1.1by Krumm [8] that demonstrates that, conditional on knowing all rational points on a particular genus 11 curve, there are no quadratic periodic points for all but finitely many $c \in \mathbf{Q}$.

Here, we would like to consider the general case wherein the coefficients lie in a field k of characteristic 0 and the domain is extended to Galois extensions K/k. In other words, if we let $\phi_c(z) = z^2 + c$ for some $c \in k$ then we consider periodic points $z \in K$ for any Galois extension K of k.[1] In this setting, we may translate the dynamical properties of periodic points to algebraic properties in Galois theory (in a different sense than the work of Morton and Patel [10]).

Let ϕ_c^N denote the N-th iterate of ϕ_c at z, i.e. $\phi_c^N = \phi_c \circ \phi_c^{N-1}$. Then we conjecture the following relation between the action of iteration and the action of $\mathrm{Gal}(\bar{k}/k)$.

Conjecture 1.4. Let k be a field of characteristic 0 and let K/k be a nontrivial finite Galois extension. Let $N \in \mathbf{N}$ be greater than 1, let $c \in k$, and let $\phi_c(z) := z^2 + c$. For any periodic point $z \in K - k$ of ϕ_c, there is an $i \in \mathbf{N}$ and a nontrivial $\sigma \in \mathrm{Gal}(K/k)$ such that

$$\phi_c^i(z) = \sigma z.$$

Remark 1.5. For $k = \mathbf{Q}$, all cycles of period up to 5 and all known 6-cycles satisfy the above conjecture, as follows from the collection of previous work in the study of quadratic periodic points of quadratic polynomials [4, 5, 6, 12, 13, 18]. Panraksa, in particular, directly considers the conjecture restricted to the setting of $k = \mathbf{Q}$ and $N = 4$ [13].

In the quadratic number field case, Conjecture 1.4 reduces finding quadratic periodic points of period N to finding rational points on certain algebraic curves, as we will see in Section 4. In particular, we demonstrate that Conjecture 1.1 is one of the consequences of Conjecture 1.4 using the description of $\mathbf{Q}$-rational periodic points of period 5 by Flynn, Poonen, and Schaefer [6].

Theorem 1.6. *There are no rational values c such that $\phi_c(z) = z^2 + c$ has a quadratic periodic point of period 5 if Conjecture 1.4 holds.*

Remark 1.7. We also obtain an analogous result to Theorem 1.6 for period 6 by using the conditional result of Stoll [17] in Corollary 4.2.

By studying Conjecture 1.4 in the Galois theoretical context, we demonstrate that it is *almost true* in the following sense. For any positive number H, consider the rational numbers of height bounded by H

$$S(H) := \{c \in \mathbf{Q} \mid H(c) \leq h\}$$

where $H(c) := \max(|a|, |b|)$ when $c = \frac{a}{b}$ in reduced form. For a property P on $\mathbf{Q}$, P is said to be *true for almost all* $c \in \mathbf{Q}$ if

$$\lim_{H \to \infty} \frac{\#P(H)}{\#S(H)} = 1$$

[1]Note that linear conjugation does not affect the length of orbits, so polynomials of the form ϕ_c capture the dynamical properties of quadratic polynomials with rational coefficients.

where $P(H) := \{c \in S(H) \mid P \text{ is true}\}$.

The definition of "almost all" can be generalized for any field k with an appropriate choice of height function H (e.g. defining the height of an algebraic number as the greatest height of the coefficients of its minimal polynomial).

Theorem 1.8. *Let k be a field of characteristic 0, K/k a nontrivial finite Galois extension, and let $N \in \mathbf{N}$ be greater than 1. Then Conjecture 1.4 is true for almost all $c \in k$.*

Remark 1.9. This "almost-ness" extends to the implications of Conjecture 1.4 described in Section 4, i.e. those statements are true for almost all $c \in k$.

2. Restrictions on orbits and Galois conjugates

First, let us define periodic points in the general sense:

Definition 2.1. Let $f : \mathbf{P}^1 \to \mathbf{P}^1$ be a non-constant map over a field k. Recall that we use f^N to denote the N-th iteration of f. If some $z \in k$ satisfies $f^N(z) = z$, then z is called a *periodic point* of f of *period* N, or an *N-periodic point* of f.

Definition 2.2. Let $f : \mathbf{P}^1 \to \mathbf{P}^1$ be a non-constant map over a field k. If $z \in k$ is an N-periodic point of f, then the orbit

$$\mathcal{O} := (z,\, f(z),\, f^2(z),\, \ldots,\, f^{N-1}(z))$$

of z is called an *N-cycle* of f. If all of the $f^i(z)$ for $i \in \{0, \ldots N-1\}$ are distinct, then $\mathcal{O}$ is called an *exact N-cycle*.

Definition 2.3. The *trace* of an N-cycle $\mathcal{O}$ is defined to be the sum of its elements.

2.1. **Geometric reformulation.** Notice that for all $N \in \mathbf{N}$, the d-periodic points of ϕ_c, for all divisors d of N, satisfy the polynomial equation $\phi_c^N(z) - z = 0$. By the Möbius inversion formula, we have that

$$\phi_c^N(z) - z = \prod_{d|N} \Phi_d(z, c)$$

for

$$\Phi_d(z, c) := \prod_{m|d} (\phi_c^m(z) - z)^{\mu(d/m)} \in \mathbf{Q}(z, c),$$

and $\Phi_d(z, c)$ is called a *dynatomic polynomial* in indeterminates z and c.

With a little bit of effort one can show that $\Phi_d(z, c)$ belongs to $\mathbf{Z}[z, c]$, and that all N-periodic points of ϕ_c are roots of the polynomial equation $\Phi_N(z, c) = 0$ (but not necessarily the converse since $\Phi_N(z, c)$ might have roots with exact period smaller than N for ϕ_c).

The polynomial equation $\Phi_N(z, c) = 0$ defines an algebraic curve in the (z, c)-plane. The affine curve $C_1(N)$ defined by the zero locus of $\Phi_N(z, c)$ is known to be smooth due to Douady–Hubbard [2] and absolutely irreducible due to Bousch [1, Chapter 3, Theorem 1]. Observe that the map ϕ_c permutes N-cycles, so the map $\sigma : (z, c) \mapsto (\phi_c(z), c)$ is an automorphism of the curve $C_1(N)$, and it generates a group $\langle\sigma\rangle$ of order equal to the largest divisor d of N such that there are exact d-periodic points of ϕ_c. Take the quotient curve $C_1(N)/\langle\sigma\rangle$, and denote the normalization of the quotient curve by $C_0(N)$. Note that for a given number field K, the K-points

on $C_0(N)$ do not necessarily arise from K-points on $C_1(N)$; rather, they correspond to $\mathrm{Gal}(\bar{K}/K)$-stable σ-orbits on $C_1(N)(\bar{k})$.

Let $p : C_1(N) \to C_0(N)$ be the quotient map. Since $\sigma : C_1(N) \xrightarrow{\sim} C_1(N)$ leaves the dominant projection $\{\Phi_N = 0\} \xrightarrow{\mathrm{pr}_2} \mathbf{A}^1$ invariant, the composite map $C_1(N) \to \mathbf{A}^1$ factors uniquely through p via a map $\pi : C_0(N) \to \mathbf{A}^1$. For each $c \in \mathbf{A}^1$ and $i \in \{0, 1\}$, define $C_i(N, c)$ to be the fiber of c on $C_i(N)$. Observe that for each point x on $C_1(N, c)(F)$ for an algebraically closed field F of characteristic 0, $p(x)$ corresponds to the σ-orbit associated to x.

Remark 2.4. The study of periodic points of exact period N in a number field K (where we also allow $c \in K$) is related to the study of K-points on the curves $C_1(N)$ and $C_0(N)$: K-points on $C_1(N)$ correspond, with finitely many exceptions (due to removal of singularities), to pairs (z, ϕ_c) for a point $z \in K$ and a map $\phi_c(z) = z^2 + c$ with $c \in K$ such that z is a periodic point of period N (not necessarily exact) of ϕ_c, and K-points on $C_0(N)$ correspond, with finitely many exceptions, to pairs $(\mathcal{O}, \phi_c)$ of a $\mathrm{Gal}(\bar{K}/K)$-stable orbit $\mathcal{O}$ and a map $\phi_c(z) = z^2 + c$ with $c \in K$; obviously these include all $(\mathcal{O}, \phi_c)$ pairs where elements of $\mathcal{O}$ are contained in K, and hence contain full information about periodic points in K.

2.2. **Galois theory reformulations.** Conjecture 1.4 can be reformulated in several different ways within the context of Galois theory. In this section, we give several equivalent statements.

First, let us refine Conjecture 1.4 with the following observation on the relation between the dynamical action of ϕ_c and the Galois action of $\mathrm{Gal}(\bar{k}/k)$ on periodic points.

Proposition 2.5. *Let k be a field of characteristic 0, K a nontrivial finite Galois extension of k of degree d, and $\phi_c(z) := z^2 + c$. Let $(z_0, \ldots, z_{N-1})$ be an exact N-cycle of ϕ_c with $N > 1$. Then the evident analogue of Conjecture 1.4 for $z := z_0$ and $c \in k$ is equivalent to the existence of an $m \in \mathbf{Z}/\gcd(N, d)\mathbf{Z}$ and a non-trivial $\sigma \in \mathrm{Gal}(K/k)$ such that*

$$\phi_c^{\frac{mN}{\gcd(N,d)}}(z) = \sigma z$$

Proof. Suppose that Conjecture 1.4 is satisfied. Let $(z_0, \ldots, z_{N-1})$ denote the N-cycle containing z with $z_0 := z$ and $z_i := \phi_c^i(z)$. Then we have some nontrivial $\sigma \in \mathrm{Gal}(K/k)$ such that $\phi_c^j(z_i) = \sigma(z_i)$ for some $i, j \in \mathbf{Z}/N\mathbf{Z}$. Note that σ commutes with ϕ_c, so we may assume that $i = 0$. Since $\sigma(z_0) = z_j$ and σ commutes with ϕ_c, we have that $\sigma \equiv \phi_c^j$ on the entire cycle.

Now recall that K is Galois, so the order of the Galois group $\mathrm{Gal}(K/k)$ is d. Thus,

$$z_0 = \sigma^d(z_0) = (\phi_c^j)^d(z_0) = z_{jd}.$$

Let r be the remainder of jd modulo N. If r is nonzero, then $(z_0, \ldots, z_{r-1})$ forms a cycle of ϕ_c with length $r < N$, violating the assumption that $(z_0, \ldots, z_{N-1})$ is an exact N-cycle. Therefore, $r = 0$; i.e., N divides jd. Consequently, j is a multiple of $\frac{N}{g}$ where $g = \gcd(N, d)$. Hence, we have some $m \in \mathbf{Z}/g\mathbf{Z}$ such that $j = m \cdot \frac{N}{g} \mod N$. □

However, we will see that Conjecture 1.4 can be restated in the following ways using Galois theory.

Proposition 2.6. *Let k be a field of characteristic 0, $c \in k$, and $N \in \mathbf{N}$ such that $N > 1$. Choose $x \in C_1(N,c)(\bar{k})$ such that x is not k-rational and let $k(x)^{\mathrm{gal}}$ denote the Galois closure of $k(x)$ over $k\big(p(x)\big)$. Then the following are equivalent:*

1. *Conjecture 1.4 for k, c, N, x, and any nontrivial finite Galois extension K/k such that x is K-rational*
2. *$k\big(p(x)\big) \neq k(x)^{\mathrm{gal}}$*
3. *either $k(x) \neq k\big(p(x)\big)$ or $k(x)$ is not Galois over $k\big(p(x)\big)$.*

Proof. The equivalence of (2) and (3) is immediate by inspecting the tower $k\big(p(x)\big) \subset k(x) \subset k(x)^{\mathrm{gal}}$. Let us consider the equivalence of (1) and (2). Let $z_j := \phi_c^j(z_0)$, and let K/k be a nontrivial finite Galois extension such that x is K-rational.

Clearly, either Conjecture 1.4 is true at z_0 or else

$$\{z_0, \ldots, z_{N-1}\} \cap \{\sigma(z_0), \ldots, \sigma(z_{N-1})\} = \emptyset$$

for all nontrivial $\sigma \in \mathrm{Gal}(K/k)$. Thus, we only need to show the equivalence of the following two conditions:

1. for every nontrivial finite Galois extension K/k such that K contains $k(x)$, there is a nontrivial $\sigma \in \mathrm{Gal}(K/k)$ such that $\sigma p(x) = p(x)$
2. $k\big(p(x)\big) \neq k(x)^{\mathrm{gal}}$

Assume (1), and take $K = k(x)^{\mathrm{gal}}$. Then $\mathrm{Gal}\Big(k(x)^{\mathrm{gal}}/k\big(p(x)\big)\Big)$ is nontrivial by our given σ and so $k\big(p(x)\big) \neq k(x)^{\mathrm{gal}}$.

Assume (2). Then for any nontrivial finite Galois extension K of k containing $k(x)$, K also contains $k(x)^{\mathrm{gal}}$, so $K \neq k\big(p(x)\big)$. Thus, the subgroup $\mathrm{Gal}\big(K/k\big(p(x)\big)\big)$ contains a non-trivial element σ. Then $\sigma p(x) = p(x)$ and so we have (1). □

Finally, we would like to rephrase the conjecture in a way that will be directly useful in the "almost theorem" in the following section.

For each $x \in C_1(N,c)$, define

$$G_x := \mathrm{Gal}\Big(k(x)^{\mathrm{gal}}/k\big(p(x)\big)\Big)$$
$$I_x := \mathrm{Gal}\Big(k(x)^{\mathrm{gal}}/k\big(x\big)\Big)$$

Then notice that (2) in Proposition 2.6 says that $G_x \neq 1$ and (3) says that either $I_x \neq 1$ or $G_x \neq I_x$. Thus, we have another reformulation of Conjecture 1.4:

Corollary 2.7. *Let k be a field of characteristic 0, $c \in k$, and $N \in \mathbf{N}$ such that $N > 1$. Choose $x \in C_1(N,c)(\bar{k})$ such that x is not k-rational. Then the following are equivalent:*

- *Conjecture 1.4 for k, c, N, x, and any nontrivial finite Galois extension K/k such that x is K-rational*
- *$G_x \neq 1$*
- *either $I_x \neq 1$ or $G_x \neq I_x$.*

3. "Almost theorems" with Hilbert irreducibility

The main conjecture formulated in this paper, Conjecture 1.4, will be difficult to prove in full generality. However, we now demonstrate that it is almost true in the sense defined in Section 1.

The idea is to move to the Galois closures of the function fields of the C_i and then show that the conjecture is true via Corollary 2.7 whenever the c-fibers of $C_i(N)_k \to \mathbf{A}_k$ are irreducible. Here, we allow $c \in k$ for a field k of characteristic 0.

Note that $C_0(N)$ and $C_1(N)$ are absolutely irreducible, hence over k [1, Chapter 3, Theorem 1]. Let $F_i(N)$ be the function fields of $C_i(N)$ and let $k(c)$ be the function field of $\mathbf{A}^1$. The projections

$$C_1 \xrightarrow{\quad p \quad} C_0 \xrightarrow{\quad \pi \quad} \mathbf{A}^1 := \mathbf{A}_k$$

given in Subsection 2.1 induce the inclusions

$$k(c) \subset F_0(N) \subset F_1(N),$$

with $F_1(N)/F_0(N)$ cyclic of degree $N > 1$.

If we let $F_1(N)^{\mathrm{gal}}$ be the Galois closure of $F_1(N)$ over $k(c)$ and let $C_1(N)^{\mathrm{gal}}$ be the normalization of $\mathbf{A}^1$ in Spec $F_1(N)$ (so $C_1(N)^{\mathrm{gal}} = \mathrm{Spec}\ R$ where R is the integral closure of $k[c]$ in $F_1(N)$), then we have the projections

$$C_1(N)^{\mathrm{gal}} \longrightarrow C_1(N) \longrightarrow C_0(N) \longrightarrow \mathbf{A}^1.$$

Define $\Gamma := \mathrm{Gal}\big(F_1(N)^{\mathrm{gal}}/k(c)\big)$. Then Γ acts on $C_1(N)^{\mathrm{gal}}$ over $\mathbf{A}^1$ and $C_1(N)^{\mathrm{gal}}/\Gamma = \mathbf{A}^1$. Away from a Γ-stable finite set of closed points in $C_1(N)^{\mathrm{gal}}$, the Γ-action on $C_1(N)^{\mathrm{gal}}$ is free and thus Γ has a free and transitive action on the fiber $C_1(N,c)^{\mathrm{gal}}$ of c in $C_1(N)^{\mathrm{gal}}$. Notice that $\mathrm{Gal}(\bar{k}/k)$ acts on $C_1(N,c)^{\mathrm{gal}}(\bar{k})$ and this action is transitive if and only if the c-fiber $C_1(N,c)^{\mathrm{gal}}$ is irreducible.

Lemma 3.1. *Assume k is a number field. The c-fiber $C_1(N,c)^{\mathrm{gal}}$ is irreducible for almost all $c \in k$.*

Proof. By Hilbert's irreducibility theorem (specifically the form on specializations of Galois groups [15, Proposition 1]) there is a thin set $\Sigma \subset k$ such that $C_1(N,c)^{\mathrm{gal}}$ consists of a single orbit under the action of $\mathrm{Gal}(\bar{k}/k)$ for all $c \notin \Sigma$. Thus, $C_1(N,c)^{\mathrm{gal}}$ is irreducible for almost all $c \in k$. □

Now using this irreducibility statement, we may prove that the main conjecture is almost true. In fact, we see that this method of proof fails in full generality only where we do not know if $C_1(N,c)^{\mathrm{gal}}$ is irreducible.

Theorem 3.2 (Restatement of Theorem 1.8)**.** *Let k be a field of characteristic 0, K/k a nontrivial finite Galois extension, and $N \in \mathbf{N}$ be greater than 1. Then Conjecture 1.4 is true for almost all $c \in k$.*

Proof. Consider c for which $C_1(N,c)^{\mathrm{gal}})(\bar{k})$ is in the locus of free-ness for the Γ-action, so $C_1(N,c)^{\mathrm{gal}}(\bar{k})$ has simply transitive Γ-action. This holds for all but finitely many c.

We can notice that the action Γ on $C_1(N,c)^{\mathrm{gal}}(\bar{k})$ commutes with the action of $\mathrm{Gal}(\bar{k}/k)$. Then the action of $\mathrm{Gal}(\bar{k}/k)$ on $C_1(N,c)^{\mathrm{gal}}$ is given by a homomorphism

$$\pi_c : \mathrm{Gal}(\bar{k}/k) \to \Gamma.$$

By Lemma 3.1, $C_1(N,c)^{\mathrm{gal}}$ is irreducible for almost all $c \in k$. This implies that π_c is surjective for c away from a thin subset of k.

Let $G := \mathrm{Gal}\big(F_1(N)^{\mathrm{gal}}/F_0(N)\big)$, a subgroup of Γ such that $C_1(N,c)^{\mathrm{gal}}/G = C_0(N)$. Note that G admits $\mathrm{Gal}\big(F_1(N)/F_0(N)\big)$ as a quotient, so $G \neq 1$ because $N > 1$. We see that G acts freely and transitively on the fiber $C_1(N,c)^{\mathrm{gal}}(\bar{k})(y)$ of each $y \in C_0(N,c)(\bar{k})$ in $C_1(N,c)^{\mathrm{gal}}(\bar{k})$ away from a finite set. The action of G on

$C_1(N,c)^{\mathrm{gal}}(\bar{k})(y)$ commutes with the natural action of $\mathrm{Gal}\big(\bar{k}/k(y)\big)$ and so we have a homomorphism

$$\pi_y : \mathrm{Gal}\big(\bar{k}/k(y)\big) \to G.$$

Clearly, $\pi_y = \pi_c|_{\mathrm{Gal}\left(\bar{k}/k(y)\right)}$ for $c = p(y)$ since we assume $c \in k$. From the surjectivity of π_c, the action of G on $C_1(N)^{\mathrm{gal}}(\bar{k})(y)$, and how G is a subgroup of Γ, we have the surjectivity of π_y for all $y \in C_0(N,c)(\bar{k})$ for almost all $c \in k$. Thus, $C_1(N,c)^{\mathrm{gal}}(\bar{k})(y)$ is irreducible over $k(y)$ with Galois group G. But this implies $G_x = \mathrm{Gal}\Big(k(x)^{\mathrm{gal}}/k\big(p(x)\big)\Big)$ is nontrivial for all $x \in C_1(N,c)(\bar{k})$ over y and so by Corollary 2.7, x satisfies Conjecture 1.4. □

4. Implications of the conjecture

In this section, we discuss the implications of Conjecture 1.4 in the special case of quadratic number fields. In this case, K is automatically Galois, so the conjecture can be applied to all quadratic number fields. Let $t \mapsto \bar{t}$ denote the nontrivial element of $\mathrm{Gal}(K/\mathbf{Q})$, and let $g = \gcd(N,2)$, so Conjecture 1.4 implies via Proposition 2.5 that there is some $m \in \mathbf{Z}/g\mathbf{Z}$ such that $z_{m\cdot\frac{N}{g}} = \bar{z_0}$.

Remark 4.1. We obtain unconditional versions of all of the following consequences with the insertion of "for almost all $c \in \mathbf{Q}$" with the application of Theorem 1.8.

If N is odd, then $g = 1$, so we have $z_0 = \bar{z_0}$; i.e., z_0 is rational. Consequently, the entire cycle lies within $\mathbf{Q}$, so we can reduce the problem of finding quadratic N-periodic points to that of finding rational N-cycles, which is a much more approachable problem (the problem of finding rational points on curves is studied extensively in the literature, whereas that of finding points within varying quadratic fields is much less common). In particular, we obtain Theorem 1.6 since Flynn, Poonen, and Schaefer [6] showed that there are no rational 5-cycles.

On the other hand, if N is even then $g = 2$, and we have either $z_0 = \bar{z_0}$, in which case the entire cycle lies within $\mathbf{Q}$; or $z_0 = \bar{z_{\frac{N}{2}}}$, in which case $z_0 + z_{\frac{N}{2}}$ is rational, and consequently the trace $z_0 + z_1 + \cdots + z_{N-1} = \sum_{j=0}^{\frac{N}{2}-1}(z_j + z_{j+\frac{N}{2}})$ is rational. Either case, the point on $C_0(N)$ that corresponds to c and the orbit $(z_0, \ldots, z_{N-1})$ is a rational point, so the problem of finding quadratic periodic points of exact period N is reduced to that of studying rational points on $C_0(N)$.

In the case of $N = 6$, the affine rational points on $C_0(6)$ are already well-understood due to a computational search by Flynn, Poonen, and Schaefer [6, Section 10] and a conditional result by Stoll [17, Theorem 7]. The computational search by Flynn, Poonen, and Schaefer found ten affine rational points on $C_0(6)$ and suggests that they are the only affine rational points on $C_0(6)$ (finiteness is known due to the high genus of $C_0(6)$). Stoll then demonstrated, assuming the weak Birch and Swinnerton-Dyer conjecture for the Jacobian of $C_0(6)$, that these are the only points of $C_0(6)(\mathbf{Q})$.

Of the ten affine rational points on $C_0(6)$ found by Flynn, Poonen, and Schaefer [6, Section 10], five lie at infinity and therefore do not correspond to any 6-cycles of quadratic polynomials $\phi_c(z) := z^2 + c$ for any $c \in \mathbf{Q}$. Four of the affine rational points correspond to 6-cycles with elements of degree > 9. The remaining known affine rational point of $C_0(6)$ corresponds to the following explicit quadratic 6-cycle

defined over $\mathbf{Q}(\sqrt{33})$:

(1)

$$c = -\frac{71}{48},\ z_0 = -1+\frac{\sqrt{33}}{12},\ z_1 = -\frac{1}{4}-\frac{\sqrt{33}}{6},\ z_2 = -\frac{1}{2}+\frac{\sqrt{33}}{12},\ z_3 = \bar{z_0},\ z_4 = \bar{z_1},\ z_5 = \bar{z_2}.$$

Therefore, we have the following corollary.

Corollary 4.2. *Let J be the Jacobian of $C_0(6)$. If*

(i) The L-series $L(J,s)$ extends to an entire function and satisfies the standard functional equation;
(ii) The weak Birch and Swinnerton-Dyer conjecture is valid for J; and
(iii) Conjecture 1.4 holds,

then the only rational c such that $\phi_c(z) = z^2 + c$ has a quadratic 6-periodic point is $-\frac{71}{48}$, and its corresponding periodic points are $z_0, \ldots, z_5$ as defined in (1).

In summary, if Conjecture 1.4 is confirmed, then it reduces all cases of finding quadratic N-periodic points to finding rational points on $C_0(N)$, which is significantly easier than finding points in quadratic extensions. In particular, for N small where we have a good understanding of $C_0(N)$, this leads to very precise (conditional) results.

5. Acknowledgements

The author would like to express thanks to Niccolò Ronchetti for suggesting this direction of study, to Zhiming Wang for the many conversations and preliminary projects that helped shape this paper, to Brian Conrad for suggesting numerous improvements and corrections, to Shou-Wu Zhang for references on Hilbert's irreducibility theorem, and to John Doyle for pointing out a few necessary clarifications. The author was partially supported by the Office of the Vice Provost for Undergraduate Education (VPUE) of Stanford University through the 2014 Stanford Undergraduate Research Institute in Mathematics (SURIM) program.

References

[1] T. Bousch. *Sur quelques problèmes de dynamique holomorphe*. PhD thesis, Université de Paris-Sud, Centre d'Orsay, 1992.
[2] A. Douady and J.H. Hubbard. *Étude dynamique des polynômes complexes (deuxième partie)*. Number 85-04 in Publications mathématiques d'Orsay. Université de Paris-Sud, Département de Mathématique, 1985.
[3] John R. Doyle. *Dynamics of quadratic polynomials over quadratic fields*. PhD thesis, University of Georgia, 2014.
[4] John R. Doyle, Xander Faber, and David Krumm. Preperiodic points for quadratic polynomials over quadratic fields. *New York J. Math.*, 20:507–605, 2014.
[5] Timo Erkama. Periodic points of quadratic polynomials. *Bulletin of the London Mathematical Society*, 38:804–814, 10 2006.
[6] E. V. Flynn, Bjorn Poonen, and Edward F. Schaefer. Cycles of quadratic polynomials and rational points on a genus-2 curve. *Duke Math. J.*, 90(3):435–463, 1997.
[7] Benjamin Hutz and Patrick Ingram. On Poonen's conjecture concerning rational preperiodic points of quadratic maps. *Rocky Mountain J. Math.*, 43(1):193–204, 2013.
[8] David Krumm. A local-global principle in the dynamics of quadratic polynomials. *International Journal of Number Theory*, 0(0):1–33, 0.
[9] Patrick Morton. Arithmetic properties of periodic points of quadratic maps. II. *Acta Arith.*, 87(2):89–102, 1998.
[10] Patrick Morton and Pratiksha Patel. The galois theory of periodic points of polynomial maps. In *Proc. London Math. Soc. 68*, pages 225–263, 1994.

[11] Patrick Morton and Joseph H. Silverman. Rational periodic points of rational functions. *Internat. Math. Res. Notices*, 1994(2):97–110, 1994.
[12] Eugen Netto. *Vorlesungen über Algebra II*. Teubner, Leipzig, 1900.
[13] Chatchawan Panraksa. *Arithmetic dynamics of quadratic polynomials and dynamical units*. PhD thesis, University of Maryland, College Park, 2011.
[14] Bjorn Poonen. The classification of rational preperiodic points of quadratic polynomials over $\mathbf{Q}$: a refined conjecture. *Math. Z.*, 228(1):11–29, 1998.
[15] Jean-Pierre Serre. *Lectures on the Mordell-Weil theorem*. Aspects of mathematics. Vieweg and Sohn, Wiesbaden, 1990.
[16] Joseph H. Silverman. *The arithmetic of dynamical systems*, volume 241 of *Graduate Texts in Mathematics*. Springer, New York, 2007.
[17] Michael Stoll. Rational 6-cycles under iteration of quadratic polynomials. *LMS J. Comput. Math.*, 11:367–380, 2008.
[18] Ralph Walde and Paula Russo. Rational periodic points of the quadratic function $Q_c(x) = x^2 + c$. *Amer. Math. Monthly*, 101(4):318–331, 1994.
[19] Zhiming Wang. Computational programs used in this paper. `https://github.com/surim14/arith-dyn`.

(Robin Zhang) Department of Mathematics, Stanford University
E-mail address: `robinz16@stanford.edu`